

Journal Article Sample

Social Sciences: Anthropology

Title Page

Full Title of Article

Levent Atici^{1*}, Suzanne E. Pilaar Birch², Burçin Erdoğan³

Author affiliations:

¹ Department of Anthropology, University of Nevada, Las Vegas, Las Vegas, NV, USA.

² Department of Anthropology & Department of Geography, University of Georgia, Athens, GA, USA.

³ Department of Archaeology, University of Thrace, Edirne, Turkey.

Corresponding author: Levent.Atici@unlv.edu (LA)

Corresponding author ORCID ID: orcid.org/0000-0002-4929-173X

Keywords: Animal domestication, spread of agriculture, island colonization, zooarchaeology, Neolithic, Turkey

Data availability: The authors confirm that all data underlying the findings are fully available without restriction. Primary or raw data used in this study will be deposited at and fully published online in the open access, peer reviewed data publishing system Open Context (<https://opencontext.org/projects/f4ab6a28-6a06-4760-a7a5-eba51d3dc964>).

Funding: This research was made possible by grants to Levent Atici (PI) and Suzanne E. Pilaar Birch from National Geographic Society (Grant# 9674-15) (<http://www.nationalgeographic.com>), to Levent Atici from the University of Nevada, Las Vegas (<http://www.unlv.edu>), to Suzanne E. Pilaar Birch from the University of Georgia (<http://www.uga.edu>), to Burçin Erdoğan from the Turkish Ministry of Culture and Tourism (http://www.kultur.gov.tr/?_dil=2) and the University of Thrace (BAP 2014-25) (<http://www-en.trakya.edu.tr>). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

1 Abstract

2 The zooarchaeological research presented here investigates Neolithic and Chalcolithic (ca.
3 6500-5000 cal. BC) animal exploitation strategies at Uğurlu Höyük on the Turkish island of
4 Gökçeada in the northeastern Aegean Sea. Toward this end, we first discuss the results of our
5 analysis of the zooarchaeological assemblages from Uğurlu Höyük and then consider the data
6 within a wider regional explanatory framework using a diachronic approach, comparing them
7 with those from western and northwestern Anatolian sites. The first settlers of Gökçeada were
8 farmers who introduced domestic sheep, goats, cattle and pigs to the island as early as 6500 years
9 BC. Our results align well with recently published zooarchaeological data on the westward
10 spread of domestic animals across Turkey and the Neolithization of southeast Europe. Using an
11 island site as a case study, we independently confirm that the dispersal of early farming was a
12 polynucleated and multidirectional phenomenon that did not sweep across the land, replace
13 everything on its way, and deliver the same “Neolithic package” everywhere. Instead, this
14 complex process generated a diversity of human-animal interactions. Thus, studying the dispersal
15 of early farmers from southwest Asia into southeast Europe via Anatolia requires a rigorous
16 methodological approach to develop a fine-resolution picture of the variability seen in human
17 adaptations and dispersals within complex and rapidly changing environmental and cultural
18 settings. For this, the whole spectrum of human-animal interactions must be fully documented for
19 each sub-region of southwest Asia and the circum-Mediterranean.

20

21 Introduction

22 The revolutionary economic and social transformation of societies from foraging to
23 farming in Southwest Asia shortly after 10,000 calibrated years BC (BC hereafter) and the
24 subsequent spread of new genes, languages, ideologies, and domesticated cereals and livestock
25 into Europe via a process called *Neolithization* from 10,000- 7000 BC have been the subjects of
26 extensive scholarly debate since the 1970s (e.g., Ammerman & Cavalli-Sforza, 1971, 1973).
27 Various models have drawn on multiple lines of converging evidence including genetics,
28 linguistics, and archaeology to explain the global dispersal of early farming populations with
29 fully developed agropastoral lifeways from primary to secondary centers of agricultural origin
30 (e.g., Arbuckle et al., 2014; Bellwood, 2009; Borić & Price, 2013; Hofmanová et al., 2016;
31 Orton, Gaastra, & Vander Linden, 2016; Özdoğan, 2005, 2011; C. Perlès, Quiles, & Valladas,
32 2013; Pinhasi, Fort, & Ammerman, 2005; Price, 2000b; Zeder, 2008, 2015).

33 Uğurlu Höyük is a Neolithic settlement on Gökçeada (Imbros in Greek), the largest
34 Turkish island situated between Anatolia and the European continent in the Aegean Sea, and
35 currently the only site with an early Neolithic component in the eastern Aegean. Thus, with its
36 key geographical location between Southeast Europe and Southwest Asia and its early Neolithic
37 strata, the results of zooarchaeological research presented here may have implications reaching
38 beyond Anatolia and contribute to our understanding of the spread and development of
39 agricultural societies in southeast Europe in general and the eastern Aegean in particular.

40 More specifically, this paper focuses on animal exploitation strategies at Uğurlu and adds
41 new zooarchaeological data to the existing body of research on the spread of domesticated
42 animals across Neolithic western Anatolia. We address the following specific questions:

43 (1) Did the islanders have a diverse subsistence strategy, including foraging and marine

44 resource exploitation, or did they heavily rely on livestock management? How did the animal
45 economy change through time?

46 (2) How did island habitation affect animal management decisions compared to the
47 mainland Anatolia? Did the islanders manage cattle, sheep, goats, and pigs differently?

48 This paper employs an analytical approach similar to that of Arbuckle and colleagues
49 (2014) in an attempt to (1) add a new site to the ‘big data’ corpus, (2) extend the scope of that
50 database spatially to go beyond the mainland Anatolia, and (3) include an island settlement to
51 compare and contrast animal exploitation strategies between the mainland Anatolia and the island
52 of Gökçeada. Toward these goals, this paper compares the results of zooarchaeological analyses
53 at Uğurlu Höyük with those from western and northwestern Anatolian sites such as Ulucak
54 Höyük, Menteşe Höyük, Çukuriçi Höyük, Ilıpınar, Barçın Höyük, Fikirtepe, and Hoca Çeşme
55 (See Fig 1 for site locations).

56 **Fig 1. Location of the sites mentioned in the text.**

57 **Conceptual framework and theoretical background**

58 In studying the dispersal of agricultural economies from southwest Asia to southeast
59 Europe, archaeologists have used a dichotomized framework. The colonization or demic
60 diffusion model entails replacement of foragers by advancing waves of farmers (Cunliffe, 2008;
61 Deguilloux, Leahy, Pemonge, & Rottier, 2012; Catherine Perlès, 2003), whereas the indigenous
62 adoption or cultural diffusion model argues for a process of acculturation instead of endemic
63 population movement and replacement (Price, 2000b and references therein). The colonization of
64 demic diffusion model hinges on the basis of the materialistic similarity with Anatolia, the
65 general absence of Mesolithic occupation on the eastern Mediterranean islands, and clear genetic

66 presence of the descendants of Near Eastern colonists in extant European populations (e.g.,
67 Cunliffe, 2008; Deguilloux et al., 2012; Özdoğan, 2007; Catherine Perlès, 2003; Price, 2000a).
68 The proponents of the latter model place emphasis on the explicit evidence for pre-pottery
69 Neolithic with Mesolithic affinities (Price, 2000a and references therein).

70 There has been a recent movement, however, toward a consensus acknowledging the
71 complexity of the processes that spread the Neolithic across Europe. Toward this end, it is now
72 recognized that farming spread into Europe by a mixture of expansion, diffusion, and adoption as
73 the predominant mechanisms (Gkiasta, Russell, Shennan, & Steele, 2003; Özdoğan, Başgelen, &
74 Kuniholm, 2012, 2013; Catherine Perlès, 2014; Robb & Miracle, 2007; Souvatzi, 2013).
75 Özdoğan (2011, 2013), Souvatzi (2013), and Perlès (2014) concur that different regions in
76 southeast Europe followed different rates of adoption of agriculture and that multiple Neolithic
77 packages successively spread from central and northwestern Anatolia to Europe.

78 **Site description and Chronology**

79 The island of Gökçeada lies about 17 km from the Gelibolu (Gallipoli) Peninsula of the
80 Anatolian mainland, and covers an area of 289.5 square km. During the Last Glacial Maximum
81 (ca. 20.000-18.000 BC), sea levels were about 120 m lower than the present sea level (Özbek,
82 2012; Van Andel & Lianos, 1983). The site of Uğurlu Höyük is a low mound covering an area of
83 approximately 250 x 200 m on a gentle slope at the eastern foot of Mount Isa (Doğanlı) on the
84 western part of the island. The site was first discovered in 1998 and a long-term project was
85 started in the summer of 2009 by Burçin Erdoğan (Erdoğan, 2011). During the six excavation
86 seasons, six main cultural phases, designated as I-VI from top to bottom, and at least 12 layers of
87 occupation have been revealed (Erdoğan, 2016). The earliest three phases (VI-IV) date to the

88 Neolithic period. Phase III is marked by the Neolithic-Chalcolithic transition, while the
89 succeeding Phase II dates to Chalcolithic. Scattered sherds from the Early Bronze Age and
90 Medieval times have been found on the surface, Phase I. Thanks to a rigorous dating program, we
91 have a well-dated and established chronology for the cultural sequence (Table S1). The earliest
92 stratum Phase VI is dated to between 6700 and 6500, Phase V between 6500 and 6000, Phase IV
93 between 5900 and 5500, and Phase III between 5400 and 4900 BC.

94 **Zooarchaeological methodology**

95 Permission to carry out the archaeological fieldwork that yielded the datasets used in this
96 project was provided by the Turkish Ministry of Culture and Tourism. All the zooarchaeological
97 specimens involved are under the auspices of the Turkish Ministry of Culture and Tourism and
98 are permanently stored in the Uğurlu Höyük Excavation Project Dig House on the island.

99 **Recovery and sampling**

100 Despite the lack of systematic dry- or wet-screening, all the excavated sediments were
101 scrutinized to ensure full recovery of macro and microfaunal remains and to minimize the effects
102 if recovery biases. Faunal assemblages from a total of 20 archaeological contexts representing
103 strata V, IV, and III (9, 7, and 4 contexts, respectively) were sampled randomly, generating 6061
104 bone fragments. Of the three strata, Phase V has generated the largest sample (N=3967), as the
105 faunal remains were densely packed in a small area of 2 x 4 m, enabling effective hand-picking.

106 **Recording**

107 The recording protocol employed in this work entailed general documentation of the
108 entire assemblage for the purpose of characterization and included every element, element

109 portion, and nonidentified splinter recovered (N=6061). No pre-sorting was practiced and all of
110 the bones were packed and stored together in the storage area of the Uğurlu Höyük dig house.
111 Every fragment was examined first by naked eye and then with a 10-15 x hand lens under strong
112 light, if necessary, for bone surface modifications, while sub-samples were randomly chosen for
113 recording variables such as fracture platform angle and percussion and notches. All the fragments
114 were identified to the maximum degree possible, refitted and mended when possible, weighed,
115 counted, labeled, assigned unique individual specimen numbers, measured when appropriate, and
116 entered into an automated FileMaker database (Levent Atici, 2011). Recording took place at the
117 project's facilities near the site on the island during field seasons 2011, 2013, and 2014 by Levent
118 Atici, and in 2015 by Levent Atici and Suzanne Pilaar Birch.

119 **Identification**

120 Taxonomic and skeletal element identifications were carried out partly using a modern
121 comparative reference collection assembled by the authors and partly using published manuals
122 describing identification criteria. When the degree of certainty of identification was high,
123 specimens were identified to the highest taxonomic category possible, i.e., species. When
124 identification to a higher taxonomic category such as species, genus, or family was not possible,
125 methodological categories, such as “medium artiodactyl” were used.

126 **Quantification**

127 Number of Fragments (NF), Number of Identified Specimens (NISP), Minimum Number
128 of Elements (MNE), and Bone Weight (BW) were quantitative measures employed in this paper
129 (Lyman, 2008). NF was used to document entire assemblages including non-specific skeletal part
130 categories and NISP was used when fragments could be identified to skeletal element and at least

131 to a taxonomic or size category (Lyman, 1994, 2008). For MNE, a combination of discrete
132 landmarks (Morlan, 1994) and manual overlap approach (Bunn & Kroll, 1986) were used.

133 Following the age data, we present biometric data from Uğurlu Höyük following the
134 standards (i.e., von den Driesch, 1976). We compare data from multiple western Anatolian
135 Neolithic sites using primary data or raw measurements directly taken from the open access, peer
136 reviewed data publishing system Open Context (<http://opencontext.org>), and/or the Logarithmic
137 Size Index (LSI) values following Richard Meadow (1999). All the datasets used in this paper
138 have citable DOIs/persistent identifiers that are listed in the appropriate supporting data tables
139 and cited accordingly in the bibliography (Levent Atici, Released 2013-02-26, Released 2013-
140 03-02; Buitenhuis, Released 2013-08-17; Canan Çakırlar, Released 2013-08-16; Galik, Released
141 2013-06-04a, Released 2013-06-04b; Gourichon & Helmer, Released 2014-05-12).

142 **Results**

143 **Assemblage formation**

144 Table S2 presents the general characteristics of the assemblages. The first step of the
145 analysis reveals the taphonomic history. Bone surface modification analysis systematically
146 included scrutiny of all skeletal parts for traces of carnivore gnawing, acid corrosion, and marks
147 left by rodents, weathering, and root etching. The analysis of 6061 fragments weighing about 26
148 kg suggests that faunal assemblages from the three strata were all accumulated, modified, and
149 destroyed largely by cultural processes.

150 A detailed analysis of bone surface modifications has revealed that rodent marks,
151 weathering, and traces of root etching are extremely rare, indicating rapid burial events and
152 intensive occupation and maintenance activities at the site. Direct and indirect traces of carnivore

153 ravaging are almost absent from the Neolithic strata (V and IV), while the Neolithic-Chalcolithic
154 transition phase (III) shows slightly increased carnivore activity at the site. The marginal number
155 (N=5) of red fox bones from Phase V and a single dog bone from Phase III independently support
156 the lack of carnivore involvement in the assemblage formation processes and can partially help
157 account for the lack of their impact as a taphonomic agent. The lack of carnivore impact, in turn,
158 indicates human processing as the primary taphonomic filter.

159 **Taxonomic composition and species trends**

160 Table S3 elaborates taxonomic composition and relative abundance of taxa based on NF,
161 MNE, and BW counts. The Uğurlu Höyük assemblages reveal that the Neolithic and Chalcolithic
162 inhabitants of the island exploited a wide range of taxa in varying proportions. The remains of
163 bovids dominate the entire cultural sequence, whereas specimens representing suids, cervids,
164 leporids, carnivores, and avifauna are present in varying and insignificant proportions and are not
165 ubiquitous. Hunted or wild taxa include large-bodied (red deer, fallow deer, and wild boar) and
166 small game (European hare). Most of the game animals identified at Uğurlu Höyük come from
167 the Neolithic strata, with level V yielding a majority of this subset. The wild cat, great bustard,
168 and mackerel shark are each represented by a single specimen from stratum IV, whereas a
169 duck/goose specimen from stratum V and a dog specimen from stratum III account for other one-
170 of-a-kind ecofacts from Uğurlu Höyük.

171 The faunal assemblages from Uğurlu are dominated by three principal food animals—
172 sheep, goats, and cattle—as their bones comprise ca. 95% of the Neolithic and 90% of the
173 Neolithic-Chalcolithic strata (Fig 2). Among the three livestock species, caprines seem to be the
174 primary focus of pastoral economy when NF and MNE counts are taken into account, as they are
175 represented in a much higher proportion (varying from 75 to 83% of all the identified bones) than

176 cattle are (varying from 10 to 20% of all the identified bones). When the bone weight data
177 presented in Table S3 are taken into account, however, the patterning changes in favor of cattle,
178 which provide the largest dietary contributions varying from 30 to 53%. Sheep outnumber goats
179 throughout the sequence, although the latter progressively increase from 6% in stratum V to 22%
180 in stratum III, whereas the exploitation of sheep and cattle visibly decline.

181 **Fig 2. Ratio distribution of principal taxa at Uğurlu Höyük using NISP counts.**

182 **Fig 3. Ternary graph showing ratio distribution of principal taxa in western Anatolia faunal**
183 **assemblages.**

184 Ulucak VI, with strata dating to 7000-6500 BC range, represents the earliest Neolithic in
185 the northern Aegean region. As Figs 6 and 7 show, Ulucak VI has a relatively even taxonomic
186 composition compared to Öküzini V, with cattle represented by ca. 16% and pigs at about 7%,
187 which indicates a multitaxic yet monodominant assemblage (sensu L. Atici, 2014). Thus, the
188 earliest phase of Ulucak Höyük is also characterized by a specialized, caprine-focused pastoral
189 economy.

190 **Fig 4. Species trends in western Anatolian faunal assemblages (%NISP).**

191 Figs 3 and 4 demonstrate a trajectory in the Aegean region toward progressively
192 increasing taxonomic evenness during the 6500-6000 BC range. At Ulucak V, while there is a
193 slight increase in the proportion of cattle from about 16 to 18%, the sharp increase in the
194 proportion of pigs from about 7 to 19% is notable and at the expense of a similarly notable drop
195 in caprine representation. Slightly later in date, Çukuriçi VIII, too, confirms the departure from a
196 caprine-dominated pastoral economy in the Aegean region. Here, the remains of cattle and pigs
197 account for about 47% (27 and 20%, respectively) of the three-tiered animal economy. When we
198 move to the northwestern region, the three Marmara sites, Fikirtepe, Barçın Höyük, and Menteşe

199 Höyük mirror this trajectory towards increased evenness in the taxonomic composition. Here,
200 too, the departure from heavy reliance on caprine management is evident. But unlike the Western
201 Anatolian region, the focus in the Marmara region shifts to cattle, not to pigs, whose
202 representation drops back to 2%.

203 **Animal exploitation: carcass management, demography of mortality, and body size**

204 Table S5 shows that all main caprine and cattle body parts are present in the assemblages
205 in varying proportions except for the total absence of axial elements for both taxa in stratum III.
206 This could be a product of small sample size and/or density-mediated attrition targeting less
207 dense axial elements, but even so, this does not indicate any clear patterning, nor does it suggest
208 selective removal, transport or processing of carcasses to primarily focus on more nutritious and
209 meaty skeletal elements. Thus, the analysis of body part distributions indicates that full caprine
210 carcasses were accessed, processed, and consumed. However, small sample sizes and disparities
211 among MNE counts do not permit meaningful body part ratio comparisons between caprines and
212 cattle, pigs, wild boars, fallow deer, and red deer (Table S5).

213 With this caveat in mind, the frequency distribution of game contrasts with that of domesticates.
214 Stratum V, with the highest NF (3,967) and MNE (954) counts among the three strata, may
215 provide the most representative picture of body part distribution for game taxa. Here, the
216 elements of forelimb and hind limb comprise 71% of all boar bones, 73% of all fallow deer
217 bones, and 50% of all red deer bones, while the elements of cranial and axial skeletons are either
218 completely absent or significantly underrepresented. Though a smaller sample, Stratum IV, too,
219 mimics the same pattern with the forelimb and hind limb elements comprising 100% of all boar
220 bones, 100% of all hare bones, and the forelimb elements making up 80% of all red deer bones.

221 For cattle, the small sample size (N=87) imposed a cut-off point and permitted the

222 assignment of cattle long bone epiphyseal specimens into either younger or older than 24 month
223 age categories. The analysis of available epiphyseal fusion data for the small sample indicates
224 that less than 30% of cattle survived beyond two years of age during stratum V with an upsurge
225 in age at death to 70% and 50% during the succeeding strata IV and III, respectively. This may
226 be due to the changing role of cattle in subsistence economy and a shift from a primary to
227 secondary animal product-oriented pastoral economy with the institutionalization and
228 intensification of farming during the late Neolithic and early Chalcolithic.

229 Although mean sheep LSI values from different Anatolian sub-regions vary
230 conspicuously, the island populations from Gökçeada during the earlier two phases, V and IV,
231 seem to align well with those from Barçın Höyük VI, Çukuriçi Höyük VIII, and Ulucak Höyük
232 VIb (Fig 5; Table S6). When placed into a longer and wider spatiotemporal framework, it
233 becomes even clearer that Uğurlu Höyük sheep represent one of the more intensively managed
234 domestic phenotypes during the Neolithic. In contrast, sheep populations during the ensuing
235 transitional Chalcolithic phase, III, must have gone through a selective process locally on the
236 island that led to further size reduction to the extent that they sit at the lowest end of the size
237 distribution.

238 **Fig 5. Distribution of *Ovis* mean LSI values for western Anatolian sites.**

239 A glance at Fig 6 (see also Table S7) reveals a similar patterning for goats with slightly
240 greater variation. Similarly, goat populations from Gökçeada fit in the range, overlapping in size
241 with other sub-regions and not representing the smallest size. Thus, it is plausible to assume that
242 Neolithic goats from Gökçeada originated from western Anatolia.

243 **Fig 6. Distribution of *Capra* mean LSI values for western Anatolian sites.**

244 For cattle, two proximal metacarpus III + IV breadth measurements, one from stratum V

245 and one from stratum IV, provide us with a glimpse into the *Bos* size range across western
246 Anatolian sites and where Uğurlu Höyük specimens fall within that range. Although neither
247 significant nor conclusive, the two specimens from Uğurlu Höyük are rather large, implying the
248 presence of either large domestic males or aurochs transported from the mainland (Fig 7).

249 **Fig 7. *Bos* spp. size distribution based on the measurement of proximal breadth (BP in mm)**
250 **in metacarpus III + IV.**

251 The biometric data presented here for Suidae are rather complicated and must be
252 interpreted with caution. On the basis of the mean LSI distributions presented in Fig 8 (see also
253 Table S8), it is hard to accurately discriminate between wild boars and domestic pigs, since
254 Epipaleolithic Öküzini V and the Cypriot Pre-Pottery Klimonas data attest to the presence of wild
255 boars whose smaller phenotypes overlap with domestic pigs. The amount of variability within
256 and among populations seems pronounced and the degree of overlap between wild boar and
257 domestic pig sizes is large. Based on the LSI patterning, we would postulate that phenotypically
258 wild and large hunted boar populations appear in the assemblages from the Marmara region: at
259 the earliest level of Ilıpınar (X), early level of Menteşe Höyük, Barçın Höyük, and Fikirtepe. In
260 contrast, all the other sub-regions indicate managed domestic pig populations. This patterning,
261 however, would be an artifact of pooling all the measurements from multiple elements to
262 overcome sample size-related biases at the expense of losing resolution. Alternatively, the
263 presence of very large male phenotypes and female-focused hunting strategies may converge to
264 skew the size distribution and make the wild, smaller female individuals fall in the domestic end
265 of the continuum. In this case, a closer look at the osteometric analysis of a single element such
266 as astragalus, which is shown in the box & whisker plot in Fig 9, could be useful. The plot shows
267 suid astragali identified as domestic, wild, and domestic or wild from Ulucak Höyük, Çukuriçi

268 Höyük, Ilıpınar, and Uğurlu Höyük. Data from the Aceramic Neolithic Klimonas from the island
 269 of Cyprus (Vigne et al., 2012) are also included to present an island wild boar population as a
 270 comparative reference. We must emphasize that the range of size distribution in domestic pigs at
 271 Ilıpınar covers domestic pigs from Ulucak Höyük and Çukuriçi Höyük and wild populations from
 272 the Cypriot Pre-Pottery Neolithic site of Klimonas and both domestic and wild individuals from
 273 Uğurlu Höyük. Therefore, the degree of overlap between the wild and domestic populations
 274 presented in the plot confirms that the biometric data are indeed nuanced, calling for careful
 275 interpretations.

276 **Fig 8. Distribution of *Sus* mean LSI values for western Anatolian sites.**

277 **Fig 9. *Sus* spp. size distribution based on the measurement of greatest lateral length (GLL in
 278 mm) in astragalus.**

279 **Concluding discussion**

280 The zooarchaeological research presented here has addressed the following specific
 281 questions to probe animal exploitation strategies at Uğurlu and to add new data to research in the
 282 spread of domesticated animals across Neolithic western Anatolia:

283 *1. Did the islanders have a diverse subsistence strategy, including foraging and marine
 284 resource exploitation, or did they heavily rely on livestock management? How did the animal
 285 economy change through time*

286 Although the Neolithic and Chalcolithic inhabitants of Gökçeada exploited a wide range
 287 of taxa in varying proportions, remains of three principal food animals—sheep, goats, and
 288 cattle—dominate the three Uğurlu Höyük assemblages. Of the taxa, caprines in general and sheep
 289 in particular were the primary focus of pastoral economy throughout the cultural sequence. Sheep

290 outnumber goats in all phases although the latter progressively increase and the exploitation of
291 sheep and cattle visibly decline by Chalcolithic.

292 During the earliest phase of the Neolithic between 7000 and 6500 BC, a more specialized,
293 caprine-dependent animal management regime seems to be represented by both sides of the
294 Aegean Sea; on the mainland Anatolia as documented at Ulucak Höyük VI and Öküzini Cave V.

295 Between 6500 and 6000 BC, Gökçeada (Uğurlu V) had a three-tiered pastoral economy
296 with a primary focus on caprines and a secondary focus on cattle; pig exploitation was marginal
297 with a proportion around 2%. In contrast, a four-tiered pastoral economy with a primary focus on
298 caprines and secondary, dual focus on cattle and pigs characterizes Çukuriçi VIII and Ulucak
299 Höyük V in the western region. Here, the ratio of pigs increases sharply as a part of progressively
300 increasing taxonomic evenness. A three-tiered animal management system with an equal focus
301 on caprines and cattle, or a shifting primary focus on either caprines or cattle is evident in the
302 Marmara and Turkish Thrace, two sub-regions of northwestern Anatolia, as documented at
303 Fikirtepe, Barçın Höyük VI, Menteşe Höyük early and late levels from the former and Hoca
304 Çeşme from the latter. The suids are represented in marginal proportions in both sub-regions.

305 During the latest phase of the Neolithic, between 6000-5500 BC, the species trend in the
306 western region shows a conspicuous continuity with a four-tiered animal husbandry, whereas the
307 sites in the Marmara Region show a greater taxonomic diversity with a sharp drop in cattle and
308 increase in caprine exploitation. The fluctuations in the reconfiguration of taxa in each region and
309 sub-region of western Anatolia mark changing roles of the four vital livestock species through
310 time and across space. This, in turn, may reflect the transformation of Neolithic societies and
311 their agropastoral economies following multiple pathways within a rapidly changing physical and
312 sociopolitical world. As far as the changes identified at Uğurlu Höyük (IV) are concerned, slight

313 but progressive increase in the exploitation of goats and decrease of sheep and cattle most likely
314 reflect the realities of resource management and impacts of environmental circumscription on an
315 island setting. Factors such as mobility, transhumance, and penning, as well as availability,
316 accessibility, predictability, and quality of grazing pastures, water, and fodder must have
317 determined animal management strategies that seem to have varied across taxa. For instance,
318 spatial constraints of islands and resource availability and abundance may pose challenges when
319 herding cattle.

320 *2. How did island habitation affect animal management decisions compared to the mainland*
321 *Anatolia? Did the islanders manage cattle, sheep, goats, and pigs differently?*

322 The analysis of body part distribution reveals nuanced and complicated data that need to
323 be interpreted cautiously. Due to sample size-related analytical biases, it is not possible to present
324 a diachronic analysis of carcass management for each livestock and game species. Still, with a
325 closer look at the earliest phase of Neolithic, Uğurlu Höyük V, somewhat representative
326 interpretations can be inferred.

327 Based on the archaeologically documented material exchanges between early farming
328 populations, it is plausible to hypothesize a process in which animals and their parts and products
329 were traded for goods among early farmers across western Anatolia. To further complicate the
330 matter, as archaeologically documented for pigs, hundreds of years of introgression between
331 feralized domestic stock and wild herds would manifest itself in the form of variable mix of traits
332 and sizes (Rowley-Conwy & Zeder, 2014: 836). This, in turn, further exacerbates the situation,
333 since a mixture of wild and domestic genetic and morphological characteristics would be
334 osteologically reflected in the zooarchaeological record. As Albarella, Dobney, and Rowley-
335 Conwy (2009) have documented, using biometry alone to accurately discriminate between wild

336 and domestic forms will not generate comparable and consistent results due to population-
337 specific intra-species size variation (see also Rowley-Conwy & Zeder, 2014: 837). Albarella and
338 colleagues (2006) note that in the islands of Corsica and Sardinia wild, feral, free-range and fully
339 domestic pigs interbreed regularly and thus create a biological continuum that could not possibly
340 be identified morphologically or biometrically, but behaviorally. As such, they treat all
341 specimens from the family Suidae as a single biological entity without attempting to assign them
342 “wild” or “domestic” status (U. Albarella et al., 2006: 292). In addition, application of multiple
343 exploitation strategies, hunting, and seasonal mobility and transhumance, may lead to distorted
344 biometric and demographic patterning that further complicates our understanding of Neolithic
345 animal management systems and obscure zooarchaeological signatures (Arbuckle & Atici, 2013).

346 The clarification of the family Suidae’s status on the island of Gökçeada and particularly
347 the verification of the presence of domestic pigs may potentially shed new light on the timing and
348 directionality of the dispersing farming populations. All four livestock species, including
349 domestic pigs with distinctively small phenotypes, are documented in the Aegean region at
350 Ulucak VI during the early seventh millennium BC, alluding to a rapid westward movement of
351 domestic animals across southern Turkey following a coastal route by sea or land (Arbuckle et
352 al., 2014).

353 Arbuckle and colleagues (2014 :8) further argue for the presence of two distinct
354 colonization pathways corresponding with distinctive animal economies and ceramic technology:
355 1) caprines, cattle, and pigs and the initial Aceramic expansion of Neolithic lifeways and with
356 later Red Slipped Burnished Ware horizon during the late eight-seventh millennium BC into
357 coastal and inland SW and western Turkey; and 2) domestic caprines and cattle associated with
358 Dark Faced Burnished Ware tradition from the interior Anatolian Plateau. Thus, would the

359 presence of domestic pigs alone place Uğurlu Höyük within the first colonization pathway and
360 directly link it to southwest and western Anatolian domain? Or would the absence of domestic
361
362 pigs suffice to establish spatiotemporal relationships between the Marmara and Thrace regions
363 and Gökçeada? The answers to these questions are nuanced and would have to incorporate more
364 than presence or absence of taxa and/or ceramic techno-typology.

365 Domestication of animals is a complex phenomenon that involves a continuum between
366 resource management, domestication or morphological changes associated with management,
367 and fully-developed animal husbandry or intentional and intensive human management of
368 animals (e.g., Arbuckle, 2013; Zeder, 2015). The study of this phenomenon, in turn, requires
369 approaches beyond binary status assignment and using single lines of evidence and/or
370 monocausal explanatory frameworks. It is difficult to clearly establish domestic status when a full
371 suite of morphological and genetic characteristics is unavailable. In the same vein, studying the
372 dispersal of early farmers from southwest Asia into southeast Europe via Anatolia requires a
373 rigorous methodological approach to develop a fine-resolution picture of the variability seen in
374 human adaptations and dispersals within complex and rapidly changing environmental and
375 cultural settings. For this, the whole spectrum of human-animal interactions must be fully
376 documented for each sub-region of southwest Asia and circum-Mediterranean. Building upon and
377 adding to the high-resolution regional-scale project spearheaded by Arbuckle and colleagues
378 (2014) to document the westward spread of domestic animals across Neolithic Turkey, Uğurlu
379 Höyük on the island of Gökçeada in the northeastern corner of the Aegean Sea, an area
380 previously under investigated and neglected, offers us an additional piece of evidence and new
381 data elaborating the nature of the Neolithic dispersals.

382 The results of zooarchaeological research presented here align well with the findings of

383 Arbuckle and Atici (2013) and Arbuckle and colleagues (2014) in that the initial diversity in
384 animal management systems of the Pleistocene-Holocene transition in southwest Asia continued
385 deep into the Neolithic and Chalcolithic with the dispersal of fully developed agropastoral
386 lifeways of early farming populations into southeast Europe. The first settlers of Gökçeada were
387 agriculturalists and they introduced domestic sheep, goats, cattle and pigs to the island as early as
388 6500 years BC. The early Neolithic has signs of continuity, but the cultures of island and
389 mainland clearly diverge. Differences in material culture may be deliberate expressions of local
390 identities within a wider cultural setting.

391 **Supporting information**

392 **Table S1.** Radiocarbon dates from Uğurlu Höyük with lab references numbers, sample numbers,
393 materials dated, and BC calibration limits for one standard error (XLSX).

394 **Table S2.** General characteristics detailing the taphonomic histories of the three Uğurlu Höyük
395 assemblages (XLSX).

396 **Table S3.** Taxonomic composition in the three Uğurlu Höyük assemblages using Number of
397 Fragments, Minimum Number of Elements, and Bone Weight in grams (XLSX).

398 **Table S4.** List of sites used in this paper with data including region, phase, chronology, author,
399 and relative abundance of *Ovis*, *Capra*, *Bos* and *Sus* based on %NISP (After Arbuckle et al. 2014)
400 (XLSX).

401 **Table S5.** Frequency distributions of body parts based on %MNE counts in main taxa (XLSX).

402 **Table S6.** Mean LSI values and standard deviations for sheep from late Pleistocene-early
403 Holocene sites in western Anatolia. Author information and links to online databases are also
404 included (XLSX).

405 **Table S7.** Mean LSI values and standard deviations for goats from late Pleistocene-early
 406 Holocene sites in western Anatolia. Links to online databases are also included (XLSX).

407 **Table S8.** Mean LSI values and standard deviations for *Sus* from late Pleistocene-early Holocene
 408 sites in western Anatolia. Author information are also included (XLSX).

409 **List of figures**

410 **Fig 1.** Location of the sites mentioned in the text: 1=Uğurlu Höyük, 2=Hoca Çeşme, 3=Yeni
 411 Kapı, 4=Fikirtepe, 5=Ilıpınar, 6=Menteşe Höyük, 7=Barçın Höyük, 8=Orman Fidanlığı,
 412 9=Ulucak Höyük, 10=Çukuriçi Höyük, and 11=Karain B and Öküzini caves.

413 **Fig 2.** Ratio distribution of principal taxa at Uğurlu Höyük using NISP counts.

414 **Fig 3.** Ternary graph showing ratio distribution of principal taxa in western Anatolia faunal
 415 assemblages. Assemblages represented are as follows: OK5=Öküzini Cave V; UG 5, 4, 3=
 416 Uğurlu Höyük V, IV, III, respectively; UL 6, 5, 4=Ulucak Höyük VI, V, IV, respectively; IPX=
 417 Ilıpınar X; HC= Hoca Çeşme; MHE=Menteşe Höyük Early; MHM=Menteşe Höyük Middle;
 418 MHL=Menteşe Höyük Late; BH6=Barçın Höyük VI; CH8=Çukuriçi Höyük VIII; FT=Fikirtepe.

419 **Fig 4.** Species trends in western Anatolian faunal assemblages (%NISP).

420 **Fig 5.** Distribution of *Ovis* mean LSI values for western Anatolian sites.

421 **Fig 6.** Distribution of *Capra* mean LSI values for western Anatolian sites.

422 **Fig 7.** *Bos* spp. size distribution based on the measurement of proximal breadth (BP in mm) in
 423 metacarpus III + IV.

424 **Fig 8.** Distribution of *Sus* mean LSI values for western Anatolian sites.

425 **Fig 9.** *Sus* spp. size distribution based on the measurement of greatest lateral length (GLL in mm)
 426 in astragalus.

427 Acknowledgements

428 We first would like to thank the Turkish Ministry of Culture and Tourism for permission
429 to conduct the archaeological fieldworks that yielded the datasets used in this project. We
430 gratefully acknowledge all the members of the Uğurlu Höyük Excavation Project for their hard
431 work which made this project possible. We extend a special thank you to Nejat Yücel whose
432 generous assistance facilitated our fieldwork. We also acknowledge and thank Burcu Birinci for
433 providing us with preliminary taxonomic identification for the shells. Finally, we thank three
434 anonymous reviewers whose insightful comments improved the paper.

435 Author Contributions

436 **Conceptualization:** Levent Atici, Suzanne E. Pilaar Birch, Burçin Erdoğan.

437 **Data curation:** Levent Atici.

438 **Formal analysis:** Levent Atici.

439 **Funding acquisition:** Levent Atici, Suzanne E. Pilaar Birch, Burçin Erdoğan.

440 **Investigation:** Levent Atici, Suzanne E. Pilaar Birch, Burçin Erdoğan.

441 **Methodology:** Levent Atici.

442 **Project administration:** Levent Atici.

443 **Resources:** Burçin Erdoğan.

444 **Writing ± original draft:** Levent Atici.

445 **Writing ± review & editing:** Levent Atici, Suzanne E. Pilaar Birch, Burçin Erdoğan.

446

447 **References**

- 448 Albarella, U., Dobney, K., & Rowley-Conwy, P. (2009). Size and shape of the Eurasian wild
449 boar (*Sus scrofa*), with a view to the reconstruction of its Holocene history.
450 *Environmental Archaeology*, 14(2), 103-136.
- 451 Albarella, U., Manconi, F., Rowley-Conwy, P., & Vigne, J.-D. (2006). Pigs of Corsica and
452 Sardinia: a biometrical re-evaluation of their status and history. In U. Tecchiati & B. Sala
453 (Eds.), *Archaeozoological studies in honour of Alfredo Riedel* (pp. 285-302). Bolzano:
454 Province of Bolzano.
- 455 Ammerman, A. J., & Cavalli-Sforza, L. L. (1971). Measuring the Rate of Spread of Early
456 Farming in Europe. *Man*, 6, 674-688.
- 457 Ammerman, A. J., & Cavalli-Sforza, L. L. (1973). A Population Model for the Diffusion of Early
458 Farming in Europe. In C. Renfrew (Ed.), *The Explanation of Culture Change* (pp. 343-
459 357). London: Duckworth.
- 460 Arbuckle, B. S. (2013). The late adoption of cattle and pig husbandry in Neolithic Central
461 Turkey. *Journal of Archaeological Science*, 40, 1805-1815.
- 462 Arbuckle, B. S., & Atici, L. (2013). Initial diversity in sheep and goat management in Neolithic
463 Southwestern Asia. *Levant*, 45(2), 219-235.
- 464 Arbuckle, B. S., Kansa, S. W., Kansa, E., Orton, D., Çakırlar, C., Gourichon, L., . . .
465 Württemberg, D. (2014). Data Sharing Reveals Complexity in the Westward Spread of
466 Domestic Animals across Neolithic Turkey. *PLoS One*, 9(6), e99845.
467 doi:10.1371/journal.pone.0099845. doi:doi:10.1371/journal.pone.0099845
- 468 Atici, L. (2009a). Implications of Age Structures for Epipaleolithic Hunting Strategies in the

- 469 Western Taurus Mountains, Southwest Turkey. *Anthropozoologica*, 44(1), 13-39.
- 470 Atici, L. (2009b). Specialisation & diversification: animal exploitation strategies in the terminal
471 Pleistocene, Mediterranean Turkey. *Before Farming*, 3, 136-152.
- 472 Atici, L. (2011). *Before the Revolution: Epipaleolithic Subsistence in the Western Taurus*
473 *Mountains, Turkey. British Archaeological Reports International Monograph Series*
474 *2251*. Oxford: Archaeopress.
- 475 Atici, L. (2014). Commingled Bone Assemblages: Insights from Zooarchaeology and
476 Taphonomy of a Bonebed at Karain B Cave, SW Turkey. In A. Osterholtz, K. Baustian, &
477 D. Martin (Eds.), *Commingled and Disarticulated Human Remains: Working Toward*
478 *Improved Theory, Method, and Data* (pp. 213-254). New York: Springer.
- 479 Atici, L. (Released 2013-02-26). Zooarchaeology of Karain Cave B: (Overview). Levent Atici
480 (Ed.) from Open Context. <[http://opencontext.org/projects/731B0670-CE2A-414A-
481 8EF6-9C050A1C60F5](http://opencontext.org/projects/731B0670-CE2A-414A-8EF6-9C050A1C60F5)> DOI:10.6078/M7CC0XMT
- 482 Atici, L. (Released 2013-03-02). Zooarchaeology of Öküzini Cave: (Overview). Levent Atici
483 (Ed.) from Open Context. <[http://opencontext.org/projects/8894EEC0-DC96-4304-
484 1EF6-4572FD91717A](http://opencontext.org/projects/8894EEC0-DC96-4304-1EF6-4572FD91717A)> DOI:10.6078/M73X84KX
- 485 Bellwood, P. (2009). The Dispersal of Established Food-Producing Populations. *Current*
486 *Anthropology*, 50(5), 621-626.
- 487
- 488 Borić, D., & Price, T. D. (2013). Strontium isotopes document greater human mobility at the start
489 of the Balkan Neolithic. *PNAS* 110(9), 3298-3303.
- 490 Buitenhuis, H. (Released 2013-08-17). Ilipinar Zooarchaeology: (Main Zooarchaeological

- 491 Dataset). Hijke Buitenhuis (Ed.). from Open Context
492 <<http://opencontext.org/tables/3746e117f8bd3c648568cdd59d00272c>> DOI
493 10.6078/M72R3PM2
- 494 Bunn, H. T., & Kroll, E. M. (1986). Systematic butchery by Plio-Pleistocene hominids at Olduvai
495 Gorge, Tanzania. *Current Anthropology*, 27(5), 431-442.
- 496 Çakırlar, C. (2012). The evolution of animal husbandry in Neolithic central-west Anatolia: the
497 zooarchaeological record from Ulucak Höyük (c. 7040–5660 cal. BC, Izmir, Turkey).
498 *Anatolian Studies*, 62(1), 1-33. doi:<http://dx.doi.org/10.1017/S0066154612000014>
- 499 Çakırlar, C. (2012). Neolithic Dairy Technology at the European-Anatolian Frontier:
500 Implications of Archaeozoological Evidence from Ulucak Höyük, İzmir, Turkey, ca.
501 7000– 5700 cal. BC. *Anthropozoologica*, 47(2), 79-100.
- 502 Çakırlar, C. (2013). Rethinking Neolithic subsistence at the gateway to Europe with
503 archaeozoological evidence from Istanbul. In M. Groot, D. Lentjes, & J. Zeiler (Eds.),
504 *The Environmental Archaeology of Subsistence, Specialisation and Surplus Food*
505 *Production* (pp. 59-79): Sidestone Press.
- 506 Çakırlar, C. (Released 2013-08-16). Zooarchaeology of Neolithic Ulucak: (Overview). Canan
507 Çakırlar (Ed.). from Open Context. <<http://opencontext.org/projects/99BDB878-6411-44F8-2D7B-A99384A6CA21>> DOI:10.6078/M7KS6PHV
508
- 509 Çakırlar, C., & Atici, L. (2017). Patterns of animal exploitation in western Turkey from
510 Palaeolithic shellfish to Byzantine elephants. In U. Albarella, H. Russ, K. Vickers, & S.
511 Viner (Eds.), *The Oxford Handbook in Zooarchaeology* (pp. 266-279). Oxford: Oxford
512 University Press.

- 513 Cannon, M. (2012). NISP, Bone Fragmentation, and the Measurement of Taxonomic Abundance.
514 *Journal of Archaeological Method and Theory*, 1-23. doi:10.1007/s10816-012-9166-z
- 515 Chalkioti, A. (2016). Reconstructing the coastal configuration of Lemnos Island (North Aegean
516 Sea, Greece) since the Last Glacial Maximum. In M. Ghilardi (Ed.), *Géoarchéologie*
517 *des îles de Méditerranée. Geoarchaeology of the Mediterranean Islands* (pp. 109-118).
518 CNRS Editions.
- 519 Çilingiroğlu, Ç., & Çakırlar, C. (2013). Towards configuring the neolithisation of Aegean
520 Turkey. *Documenta Praehistorica*, 40, 21-29.
- 521 Cunliffe, B. (2008). *Europe Between the Oceans: 9000 bc-AD 1000*. New Haven: Yale
522 University Press.
- 523 De Cupere, B., Duru, R., & Umurtak, G. (2008). Animal husbandry at the Early Neolithic to
524 Early Bronze Age site of Bademağacı (Antalya province, SW Turkey): evidence from the
525 faunal remains. In E. Vila, L. Gourichon, A. M. Choyke, & H. Buitenhuis (Eds.),
526 *Proceedings of the 8th International Symposium on the Archaeozoology of Southwestern*
527 *Asia and Adjacent Areas* (pp. 367-405). Lyon: Archéorient, Maison de l'Orient et de la
528 Méditerranée.
- 529 Deguilloux, M.-F., Leahy, R., Pemonge, M.-H., & Rottier, S. (2012). European Neolithization
530 and Ancient DNA: An Assessment. *Evolutionary Anthropology: Issues, News, and*
531 *Reviews*, 21(1), 24-37. doi:10.1002/evan.20341
- 532 Deniz, E., & Payne, S. (1982). Eruption and wear in the mandibular dentition as a guide to
533 ageing Turkish Angora goats. In B. Wilson, C. Grigson, & S. Payne (Eds.), *Ageing and*
534 *Sexing Animal Bones from Archaeological Sites* (Vol. 109, pp. 155-205). Oxford:
535 Archaeopress.

- 536 Erdoğan, B. (2011). Preliminary Report from the 2009 and 2010 Field Seasons at Uğurlu on the
537 Island of Gökçeada. *Anatolica*, XXXVII, 45-65.
- 538 Erdoğan, B. (2013). Uğurlu: A Neolithic settlement on the Aegean Island of Gökçeada. In M.
539 Özdoğan, N. Başgelen, & P. Kuniholm (Eds.), *The Neolithic in Turkey. New Excavations*
540 *and New Research. Northwestern Turkey and Istanbul* (Vol. 5). Istanbul: Archaeology
541 and Art Publications.
- 542 Erdoğan, B. (2016). The Neolithic landscape and settlement of the Island of Gökçeada (Imbros,
543 Turkey). In M. Ghilardi (Ed.), *Géoaarchéologie des îles de Méditerranée.*
544 *Geoarchaeology of the Mediterranean Islands* (pp. 89-94): CNRS Editions.
- 545 Erdoğan, B. (2017). Whither The Aegean Neolithic. In Z. Tsirtsoni, A. Reingruber, & P.
546 Nedelcheva (Eds.), *Going West? The spread of farming between the Bosphorus and the*
547 *Lower Danube Region*. New York: Routledge.
- 548 Evershed, R. P., Payne, S., Sherratt, A. G., Copley, M. S., Coolidge, J., Urem-Kotsu, D., . . .
549 Burton, M. M. (2008). The Earliest date for milk use in the Near East and southeastern
550 Europe linked to cattle herding. *Nature*, 455, 528-531.
- 551 Galik, A. (Released 2013-06-04a). Barçın Höyük Zooarchaeology: (Overview). Alfred Galik
552 (Ed.) from Open Context. <<http://opencontext.org/projects/74749949-4FD4-4C3E-C830-5AA75703E08E>> DOI:10.6078/M78G8HM0
- 553
- 554 Galik, A. (Released 2013-06-04b). Çukuriçi Höyük Zooarchaeology: (Overview). Alfred Galik
555 (Ed.). from Open Context. <<http://opencontext.org/projects/BC90D462-6639-4087-8527-6BB9E528E07D>> DOI:10.6078/M7D798BQ
- 556
- 557 Gkiasta, M., Russell, T., Shennan, S., & Steele, J. H. (2003). Neolithic transition in Europe: the

- 558 radiocarbon record revisited. *Antiquity*, 77, 45-62.
- 559 Gourichon, L., & Helmer, D. (Released 2014-05-12). Faunal Data from Neolithic Mentese.
- 560 Lionel Gourichon and Daniel Helmer (Eds.). from Open Context.
- 561 <http://opencontext.org/projects/42EAD4DB-BAED-4A58-9B9B-7EC85266D2A9>> DOI:
- 562 <http://dx.doi.org/10.6078/M7S46PVN>
- 563 Gurova, M. (2008). Towards an understanding of Early Neolithic populations: a flint perspective
- 564 from Bulgaria. *Documenta Praehistorica XXXV*, 111-129.
- 565 Hofmanová, Z., Kreutzer, S., Hellenthal, G., Sell, C., Diekmann, Y., Díez-del-Molino, D., . . .
- 566 Burger, J. (2016). Early farmers from across Europe directly descended from Neolithic
- 567 Aegeans. *PNAS*, 113(25), 6886-6891. doi:10.1073/pnas.1523951113
- 568 Hongo, H., & Meadow, R. H. (2000). Faunal Remains from Pre-Pottery Neolithic Levels at
- 569 Cayonu, Southeastern Turkey,: A Preliminary Report Focusing on Pigs (*Sus* sp.). In M.
- 570 Mashkour, A. M. Choyke, H. Buitenhuis, & F. Poplin (Eds.), *Archaeozoology of*
- 571 *Southwest Asia* (pp. 121-140). Groningne: Oxbow Books.
- 572 Lichter, C. (2011). Neolithic Stamps and Neolithization Process. A Fresh Look at an Old Issue In
- 573 R. Krauß (Ed.), *Beginnings: New Research in the Appearance of the Neolithic between*
- 574 *Northwest Anatolia and the Carpathian Basin* (Vol. Menschen - Kulturen - Traditionen;
- 575 *Forschung Cluster 1, Band 1, pp. 33-44*): Rahden: Verlag Marie Leidorf GmbH.
- 576 Lyman, R. L. (1994). Quantitative units and terminology in zooarchaeology. *American Antiquity*,
- 577 59(1), 36-71.
- 578 Lyman, R. L. (2008). *Quantitative Paleozoology*. Cambridge: Cambridge University Press.
- 579 Meadow, R. H. (1999). The use of size index scaling techniques for research on

- 580 archaeozoological collections from the Middle east. In A. Von Den Driesch (Ed.),
581 *Historia Animalium Ex Ossibus: Beitrage zur Palaoanatomie, Archaologie, Agyptologie,*
582 *Ethnologie und Geschichte der Tiermedizin* (pp. 285-300). Rahden: Verlag Marie Leidorf
583 GmbH.
- 584 Morlan, R. E. (1994). Bison bone fragmentation and survivorship: a comparative method.
585 *Journal of Archaeological Science*, 21, 797-807.
- 586 Munro, N. D., & Stiner, M. C. (2015). Zooarchaeological Evidence for Early Neolithic
587 Colonization at Franchthi Cave (Peloponnese, Greece). *Current Anthropology*, 56(4), 596-
588 603. doi:<http://www.jstor.org/stable/10.1086/682326>
- 589 Orton, D. C., Gaastra, J., & Vander Linden, M. (2016). Between the Danube and the Deep Blue
590 Sea: zooarchaeological meta-analysis reveals variability in the spread and development of
591 Neolithic farming across the western Balkans. *OpenQuaternary*, 2, 6. doi:DOI:
592 <http://doi.org/10.5334/oq.28>
- 593 Ottoni, C., Girdland Fink, L., Evin, A., Geörg, C., De Cupere, B., Van Neer, W., . . . Larson, G.
594 (2013). Pig domestication and human-mediated dispersal in western Eurasia revealed
595 through ancient DNA and geometric morphometrics. *Molecular Biology and Evolution*,
596 30(4), 824-832.
- 597 Özbal, H., Thissen, L., Doğan, T., Gerritsen, F., Özbal, R., & Türkekul-Biyik, A. (2013). Neolitik
598 Batı Anadolu ve Marmara yerleşimleri çanak çömleklerinde organik kalıntı analizleri.
599 *Arkeometri Sonuçları Toplantısı*, 28, 105-114.
- 600 Özbek, O. (2012). Sea-level changes and Prehistoric sites on the coasts of Southern Turkish
601 Thrace, 12000-6000 BP. *Quaternary International* doi: 10.1016/j.quaint.2012.01.006.

- 602 Özdoğan, M. (2005). Westward expansion of the Neolithic way of life: what we know and what
603 we do not know. In C. Lichter (Ed.), *How did farming reach Europe? Anatolian*
604 *European relations from the second half of the 7th through the first half of the 6th*
605 *millennium cal BC. Proceeding of the International Workshop Istanbul, 20-22 May 2004*
606 (pp. 13-27). Istanbul: Ege Yayinlari.
- 607 Özdoğan, M. (2007). Coastal changes of the Black Sea and Sea of Marmara in archaeological
608 Perspective. In V. Yanko-Hombach, A. S. Gilbert, N. Panin, & P. M. Dolukhanov (Eds.),
609 *The Black Sea Flood Question: Changes in Coastline, Climate and Human Settlement*
610 (pp. 651-669). Dordrecht, The Netherlands: Springer.
- 611 Özdoğan, M. (2011). Archaeological Evidence on the Westward Expansion of Farming
612 Communities from Eastern Anatolia to the egean and the Balkans. *Current Anthropology*,
613 52(S4), S415-S-430.
- 614 Özdoğan, M. (2013). Neolithic Sites in the Marmara Region: Fikirtepe, Pendik, Yarımburgaz,
615 Toptepe, Hoca Çeşme and Aşağı Pınar. In M. Özdoğan, N. Başgelen, & P. Kuniholm
616 (Eds.), *The Neolithic in Turkey. New Excavations and New Research. Northwestern*
617 *Turkey and Istanbul* (pp. 167-269). Istanbul: Archaeology and Art Publications.
- 618 Özdoğan, M., Başgelen, N., & Kuniholm, P. (Eds.). (2012). *The Neolithic in Turkey. New*
619 *Excavations and New Research. Western Turkey* Istanbul: Archaeology and Art
620 Publications.
- 621 Özdoğan, M., Başgelen, N., & Kuniholm, P. (Eds.). (2013). *The Neolithic in Turkey. New*
622 *Excavations and New Research. Northwestern Turkey and Istanbul*. Istanbul:
623 Archaeology and Art Publications.
- 624 Perissoratis, C., & Conispoliatis, N. (2003). The impacts of sea-level changes during latest

- 625 Pleistocene and Holocene times on the morphology of the Ionian and Aegean seas (SE
626 Alpine Europe). *Marine Geology*, 196, 145-156.
- 627 Perlès, C. (2003). An alternate (and old-fashioned) view of Neolithisation in Greece. *Documenta*
628 *Praehistorica*, XXX, 99-113.
- 629 Perlès, C. (2014). Neolithic Anatolia as seen from the West. In M. Özdoğan, N. Başgelen, & P.
630 Kuniholm (Eds.), *The Neolithic in Turkey. 10500-5200 BC: Flora, Fauna, Dating,*
631 *Symbols of Belief, with Views from North, South, East, and West* (pp. 403-432). Istanbul:
632 Archaeology and Art Publications.
- 633 Perlès, C., Quiles, A., & Valladas, H. (2013). Early seventh-millennium AMS dates from
634 domestic seeds in the Initial Neolithic at Franchthi Cave (Argolid, Greece). *Antiquity*,
635 87(338), 1001–1015.
- 636 Pilaar Birch, S. (2017). From the Aegean to the Adriatic: Exploring the Earliest Neolithic Island
637 Fauna. *Journal of Island and Coastal Archaeology*, DOI:
638 10.1080/15564894.2017.1310774.
- 639 Pilaar Birch, S., Çakırlar, C., Scheu, A., & Buckley, M. (submitted). Multiproxy analyses of
640 sheep and goat remains from Neolithic Ulucak in the eastern Aegean. *Archaeological and*
641 *Anthropological Sciences*.
- 642 Pinhasi, R., Fort, J., & Ammerman, A. J. (2005). Tracing the Origin and Spread of Agriculture in
643 Europe. *PLoS Biology*, 3(12), e410. doi:410.1371/journal.pbio.0030410.
- 644 Price, T. D. (2000a). Europe's first farmers: an introduction. In T. D. Price (Ed.), *Europe's First*
645 *Farmers* (pp. 1-18). Cambridge: Cambridge University Press.
- 646 Price, T. D. (Ed.) (2000b). *Europe's First Farmers*. Cambridge: Cambridge University Press.

- 647 Reingruber, A. (2011). Early Neolithic Settlement Patterns and Exchange Network in the Aegean.
648 *Documenta Praehistorica*, 38, 291-305.
- 649 Robb, J., & Miracle, P. (2007). Beyond 'migration' versus 'acculturation': new models for the
650 spread of agriculture In A. Whittle & V. Cummings (Eds.), *Going Over: The Mesolithic-*
651 *Neolithic Transition in North-West Europe. Proceedings of the British Academy 144* (pp.
652 99-115): The British Academy.
- 653 Rowley-Conwy, P., & Zeder, M. A. (2014). Wild Boar or Domestic Pigs? Response to Evin et al.
654 *World Archaeology*, 46(5), 835-840. doi:10.1080/00438243.2014.953712
- 655 Shipman, P. (1981). *Life History of a Fossil: An Introduction to Taphonomy and Paleoecology*.
656 Cambridge: Harvard University Press.
- 657 Simmons, A. (2007). *The Neolithic Revolution in the Near East: Transforming the Human*
658 *Landscape*. Tuscon: The University of Arizona Press.
- 659 Sondaar, P. Y. (1977). Insularity and its effects on mammal evolution: 671-707. In H. M. K., G.
660 P. C., & H. B. M (Eds.), *Major Patterns in Mammal Evolution*. New York Plenum.
- 661 Souvatzi, S. (2013). Diversity, Uniformity and the Transformative Properties of the House in
662 Neolithic Greece. In D. Hofmann & J. Smyth (Eds.), *Tracking the Neolithic House in*
663 *Europe: Sedentism, Architecture and Practice* (pp. 45-64). New York: Springer.
- 664 Stiner, M. C. (1990). The use of mortality patterns in archaeological studies of hominid predatory
665 adaptations. *Journal of Anthropological Archaeology*, 9, 305-351.
- 666 Uerpmann, M., & Uerpmann, H.-P. (1994). Animal bone finds from excavation 520 at Qala'at al-
667 Bahrain. In F. Holland & H. H. Anderson (Eds.), *The Northern City Wall and the Islamic*
668 *Fortress, vol. 1 of Qala'at al-Bahrain* (pp. 417-444). Aarhus: Jutland Archaeological

- 669 Society.
- 670 Van Andel, T. H., & Lianos, N. (1983). Prehistoric and Historic shorelines of the southern Argolid
671 Peninsula: a subbottom profiler study. *Nautical Archaeology* 12(4), 303-324.
- 672 Vigne, J.-D., Briois, F., Zazzo, A., Willcox, G., Cucchi, T., Thiébaud, S., . . . Guilaine, J. (2012).
673 First wave of cultivators spread to Cyprus at least 10,600 y ago. *PNAS*, 109(22), 8445-
674 8449.
- 675 von den Driesch, A. (1976). *A Guide to the Measurement of Animal Bones from Archaeological*
676 *Sites: As Developed by the Institut Für Palaeoanatomie, Domestikationsforschung und*
677 *Geschichte Der Tiermedizin of the University of Munich*. Cambridge, MA: Peabody
678 Museum of Ethnology and Archaeology, Harvard University.
- 679 Zeder, M. A. (2006). Reconciling Rates of Long Bone Fusion and Tooth Eruption and Wear in
680 Sheep (*Ovis*) and Goat (*Capra*). In D. Ruscillo (Ed.), *Ageing and Sexing Animals from*
681 *Archaeological Sites* (pp. 87-118). Oxford: Oxbow Press.
- 682 Zeder, M. A. (2008). Domestication and early agriculture in the Mediterranean Basin: Origins,
683 diffusion, and impact. *PNAS*, 105(33), 11597-11604.
- 684 Zeder, M. A. (2015). Core Questions in Domestication Research. *PNAS*,
685 [doi/10.1073/pnas.1501711112](https://doi.org/10.1073/pnas.1501711112). doi:doi/10.1073/pnas.1501711112