

Publications

Posakony LG, Raftery LA, Gelbart WM. Wing formation in *Drosophila* requires *decapentaplegic* gene function along the anterior-posterior compartment boundary. *Mech Dev* 1991; 33:69-82.

Blackman RK, Sanicola M, Raftery LA, Gillevet T, Gelbart WM. An extensive 3' cis-regulatory region directs the imaginal disk expression of *decapentaplegic*, a member of the TGFbeta family in *Drosophila*. *Development* 1991; 111:657-65.

Raftery LA, Sanicola M, Blackman RK, Gelbart WM. The relationship of *decapentaplegic* and engrailed expression in *Drosophila* imaginal disks: do these genes mark the anterior-posterior compartment boundary? *Development* 1991; 113:27-33.

Raftery LA, Twombly V, Wharton K, Gelbart WM. Genetic screens to identify elements of the *decapentaplegic* signaling pathway in *Drosophila*. *Genetics* 1995; 139:241-54.

Sekelsky JJ, Newfeld SJ, Raftery LA, Chartoff EH, Gelbart WM. Genetic characterization and cloning of *Mothers against dpp*, a gene required for *decapentaplegic* function in *Drosophila melanogaster*. *Genetics* 1995; 139:1347-58.

Baden H, Kollias N, Anderson RR, Hopkins T, Raftery LA. *Drosophila melanogaster* larvae detect low doses of UVC radiation as manifested by a writhing response. *Arch Insect Biochem Physiol* 1996; 32:187-96.

Raftery LA, Wisotzkey RG. Characterization of *Medea*, a gene required for maximal function of the *Drosophila* BMP homolog, *decapentaplegic*. *Ann. NY Acad. Sci.* 1996; 785:318-320.

Dobens LL, Hsu T, Twombly V, Gelbart WM, Raftery LA, Kafatos FC. The *Drosophila bunched gene* is a homologue of the growth factor stimulated mammalian TSC22 sequence and is required for proper follicle cell patterning during oogenesis. *Mech Dev* 1997; 65:197-208.

Dobens L, Raftery LA. *Drosophila* oogenesis: a model system to understand TGFbeta/Dpp directed cell morphogenesis. *Ann NY Acad Sci* 1998; 857:245-247.

Wisotzkey RG, Mehra A, Sutherland D, Dobens LL, Liu X, Dohrmann C, Attisano L, Raftery LA. *Medea* is a *Drosophila SMAD4* homolog that is differentially required to potentiate DPP responses. *Development* 1998; 125:1433-144.

Brummel T, Abdollah S, Haerry TE, Shimell MJ, Merriam J, Raftery L, Wrana J, O'Connor MB. The *Drosophila* Activin receptor Baboon signals through dSmad2 and controls cell proliferation but not patterning during larval development. *Genes Dev* 1999; 13:98-111.

- Dohrmann CE, Belaoussoff M, Raftery LA. Dynamic expression of TSC-22 at sites of epithelial-mesenchymal interactions during mouse development. *Mech Dev* 1999; 84:147-151.
- Raftery, LA, Sutherland, DJ. From *Mad* to Smads: TGFb??? signal transduction in *Drosophila* development. *Dev Biol* 1999; 210:251-268.
- Dobens LL, Peterson J, Treisman J, Raftery LA. *Drosophila bunched* integrates opposing DPP and EGF signals to set the operculum boundary. *Development* 2000; 127:745-754.
- Dobens LL, Raftery LA. Integration of epithelial patterning and morphogenesis in *Drosophila* ovarian follicle cells. *Dev Dyn* 2000; 218:80-93.
- Dai H, Hogan C, Gopalkrishnan B, Torres-Vasquez J, Nguyen M., Park S, Raftery LA, Warrior R, Arora K. The zinc finger protein Schnurri acts as a Smad partner in mediating the transcriptional response to Dpp. *Dev. Biol.* 2000; 227:373-387.
- Dobens LL, Martin-Blanco E, Martinez-Arias A, Kafatos FC, Raftery LA. *Drosophila puckered* regulates Fos/Jun levels during follicle cell morphogenesis. *Development* 2001; 128:1845-1856.
- Dohrmann CE, Noramly S, Raftery LA, Morgan, B. Opposing effects on TSC-22 expression by BMP and receptor tyrosine kinase signals in the developing feather field. *Dev. Dyn.* 2002; 223:85-95.
- Gupta RA, Sarraf P, Brockman JA, Shappell SB, Willson TM, Raftery, L, DuBois, RN. Peroxisome proliferator-activated receptor γ and transforming growth factor- β pathways inhibit intestinal epithelial cell growth by regulating levels of TSC-22. *J Biol Chem* 2003; 278:7431-7438.
- Soma T, Dohrmann CE, Hibino T, Raftery LA. Profile of TGF β responses in the murine hair cycle. *J Inv Derm:* 2003; 121:969-975.
- Sutherland DJ, Li M, Liu X, Stefancsik R, Raftery LA. Stepwise formation of a Smad activity gradient during dorsal-ventral patterning of the *Drosophila* embryo. *Development* 2003; 130:5705- 5716.
- Raftery, LA, Sutherland, DJ Gradients and Thresholds: BMP response gradients unveiled in *Drosophila* embryos. *Trends Genet* 2003; 19:667-724.
- Whitman M, Raftery L TGR β at the summit. *Development* 2005; 132(19): 4205-4210.
- Dobens L, Jaeger A, Peterson JS, Raftery LA bunched sets a boundary for Notch signaling to pattern anterior eggshell structures during *Drosophila* oogenesis. *Dev. Biol.* 2005; 287:425-437.

Raftery LA, Korochkina S, Cao J. “Smads in *Drosophila* – interpretation of graded signals in vivo.” in Smad Signal Transduction, C-H Heldin, P ten Dijke, eds. Springer, 2006, 55-73.

Cao J, Pellock BJ, White K, Raftery LA. A commercial phospho-Smad antibody detects endogenous BMP signaling in *Drosophila* tissues. *Drosophila Information Service* 2006; 89:131-135.

Gluderer S, Köhler K, Oldham S, Rintelen F, Sulzer A, Schütt C, Wu X, Raftery L, Hafen E, Stocker H. Bunched, the *Drosophila* homolog of the mammalian tumor suppressor TSC-22, promotes cellular growth. *BMC Dev. Biol.* 2008 8:10.

Wu X, Tanwar PS, Raftery LA, *Drosophila* follicle cells: morphogenesis in an eggshell. *Seminars Cell Dev. Biol.* 2008; 19:271-282.

Wu X, Yamada-Mabuchi M, Morris EJ, Gluderer S, Tanwar P, Dobens L, Cao J, Stocker H, Hafen E, Dyson NJ, Raftery LA. The *Drosophila* homolog of human tumor suppressor TSC-22 promotes cellular growth, proliferation and survival. *Proc. Natl. Acad. Sci.* 2008; 105:5414-5419.

Bilenca A, Cao J, Colice M, Ozcan A, Bouma B, Raftery L, Tearney G. Fluorescence interferometry: principles and applications in biology. *Ann. N.Y. Acad. Sci.* 2008; 1130:68-77.

Batut J, Schmierer B, Howell M, Cao J, Raftery LA, Hill CS. Two highly related regulatory subunits of PP2A exert opposite effects on TGFbeta signaling. *Development* 2008; 135:2927-2937.

Miles WO, Jaffray E, Campbell SG, Takeda S, Bayston LJ, Basu SP, Li M, Raftery LA, Ashe MP, Hay RT, Ashe HL. Medea SUMOylation restricts the signaling range of the Dpp morphogen in the *Drosophila* embryo. *Genes Dev.* 2008; 22:2578-2590.

Raftery LA, Umulis, D Generation and interpretation of BMP gradients in *Drosophila*, *Cur. Opin. Cell Biol.* 2012, 24:1-8.

Brooks AI, Dou W, Pargett M, Brosnan T, Raftery LA, Umulis DM. BMP signaling in development: A critical analysis of imaging and interpretation. *FEBS Lett.* 2012, 586:1942-1952.

Peterson AJ, Jensen PA, Shimell MJ, Stefancsik R, Wijayatunge R, Herder R, Raftery LA, O'Connor MB.. R-Smad competition controls Activin receptor output in *Drosophila*. *PLoS One* 2013, 7: e36548.